

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) Publication number:

0 380 230
A2

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 90300456.2

(51) Int. Cl.⁵ **A61K 37/24, C07K 99/30**

(22) Date of filing: 17.01.90

(30) Priority: 24.01.89 US 301353
12.05.89 US 351193

(43) Date of publication of application:
01.08.90 Bulletin 90/31

(84) Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

(71) Applicant: **APHTON CORPORATION**
P.O. Box 1049
Woodland, CA 95695(US)


(72) Inventor: **Gevas, Philip C.**
487-A Portlock Road
Honolulu, Hawaii 96825(US)
Inventor: **Grimes, Stephen**
1400 Cypress Lane
Davis, California 95616(US)
Inventor: **Karr, Stephen L., Jr.**
624 Villanova Drive
Davis, California 95616(US)
Inventor: **Littenberg, Richard L.**
70 Kai One Place
Kai Lua, Hawaii 96734(US)

(74) Representative: **Lucas, Brian Ronald**
Lucas, George & Co. 135 Westhall Road
Warlingham Surrey CR3 9HJ(GB)

(54) Immunogenic compositions against gastrin peptides.

(57) Immunogenic compositions useful for the treatment of ulcers or tumors whose growth is dependent on or stimulated by gastrin hormones are disclosed. The immunogenic compositions induce antibodies in a subject which selectively neutralize the specific hormones. Pharmaceutical compositions comprising effective amounts of the immunogenic compositions and methods of treatment using the compositions are disclosed. A method of reversing the inventive treatments by neutralizing the antibodies induced in vivo is also disclosed.

Figure 3
Stomach Acid Output in Response to
Injection of G17 and Pentagastrin (pG)
in Rat Actively Immunized Against G17



Arrows: Time of G17 or pG injection.
Acid Secretion peaks are numbered 1,2,3.

Xerox Copy Centre

EP 0 380 230 A2

IMMUNOGENIC COMPOSITIONS AGAINST GASTRIN PEPTIDES

Background of the Invention

Peptic ulcer disease exists in two forms, duodenal ulcers and gastric ulcers. Central to the cause of duodenal ulcers, is the production of excess stomach acid and pepsin and a rapid gastric emptying time. This results in an increase in duodenal exposure to secreted acid and enzymes, and in mucosal damage.

The second form of the disorder, gastric ulcer disease, may be caused by increased stomach acid and a breakdown of the complex stomach defenses that normally protect the gastric mucosa from acid damage. Although the two conditions have different etiologies, both benefit from a reduction in gastric acid secretion.

Because excess stomach acid is a central cause of ulcers, antacid preparations are commonly used as one method of treatment. This method merely neutralizes stomach acid after it is produced. Consequently, large quantities of antacids must be consumed on an ongoing basis to neutralize acid which is continually produced in the stomach. Antacids do not cure the disease because they do not affect the source of acid production.

Gastric acid is produced in a specialized stomach cell, the parietal cell. Parietal cells can be stimulated to secrete acid by acetylcholine, histamine and gastrin, upon the binding of each of these compounds with specific receptors on the surface of the cell. Of these the most potent stimulator of acid secretion is the peptide hormone gastrin.

Current approaches to the control and cure of peptic ulcers center upon devising drugs that inhibit the ability of one or more of these compounds to stimulate acid production or secretion. The most effective group of drugs approved for sale are the H₂ antagonists (e.g. Tagamet® and Zantac®) which block the histamine H₂ receptors on gastric parietal cells and inhibit acid secretion. These drugs, however, require relatively large doses on a daily basis and may induce several undesirable side effects. In cases where H₂ antagonists have cured ulcers, relapses occur in almost 100% of cured individuals within a year of discontinuation of treatment. Other drugs have also exhibited problems, including low efficacy and unacceptable levels of toxicity. In the case the peptide hormone gastrin, no successful chemical antagonists have been identified.

Gastrin has several important functions in the gastrointestinal tract, the two most important being stimulation of acid secretion and stimulation of the growth of cells in the gastrointestinal tract. The hormone exists in at least two molecular forms, heptadecagastrin ("G₁₇") and tetracosagastrin ("G₃₄") named according to the number of amino acid ("AA") residues in each molecule. G₃₄ and G₁₇ are identical in structure at the carboxy terminus, which is the binding site of the hormones with receptors. G₁₇ constitutes the 17 carboxy terminal ("C-terminal") end residues of G₃₄. G₃₄ consists of the 17 C-Terminal end residues which comprise G₁₇ and an additional different amino acid sequence of 17 amino terminal ("N-terminal") residues. When G₃₄ is split by trypsin a G₁₇ subunit and a non-hormonal 17 amino acid subunit results. Though G₁₇ is usually obtained by trypsin cleavage of G₃₄, each form may also be generated separately from its own prohormone.

Although G₁₇ and G₃₄ are thought to be equipotent on a molar basis as stimulators of acid release, G₃₄ is most probably responsible for the stimulation of growth of the gastrointestinal mucosa and the maintenance of the basal acidity of the stomach. G₃₄ is the principal form present during interdigestive periods. G₃₄ has a serum half life approximately six times as long as G₁₇ (40 minutes versus 6 minutes) and is produced in both the stomach and the duodenum. Alternatively, G₁₇ is the primary stimulator of meal induced gastric acid secretion. G₁₇ is 1500 times more potent than histamine and makes up 90% of the antral (stomach) gastrin. G₁₇ accounts for roughly 60%-70% of the gastrin-mediated acid release.

The prior art in the area of gastrin immunology mainly concerns the induction of antibodies useful for identifying anatomic sites containing or producing gastrin G₁₇ or G₃₄ in laboratory animals; see Sugano, K., et al., 1985, "Identification and characterization of glycine-extended post translational processing intermediates of progastrin in porcine stomach", *J. of Biological Chemistry* 260: 11724-11729; Vaillant, C., et al., 1979, "Cellular origins of different forms of gastrin: The specific immunocytochemical localization of related peptides. *J. Histochem Cytochem* 27:932-935; Larsson, L.I. et al., 1977, "Characterization of antral gastrin cells with region-specific antisera". *J. Histochem. Cytochem* 25: 1317-1321. The antisera reported in these publications contained antibodies of numerous specificities, for a variety of antigenic epitopes on gastrin molecules.

Attempts to control gastrin levels by anti-gastrin antibodies induced by active immunization or passive administration of preformed antibodies such as those reported in Jaffe, B.M., et al., 1971, "Gastrin

resistance following immunizations to the C-terminal tetrapeptide amide of gastrin, Surgery 69: 232-238; Jaffe, B.M., et al., 1970, "Inhibition of endogenous gastrin activity by antibodies to the carboxyl terminal tetrapeptide amide of gastrin", Gastroenterology 58: 151-156; Jaffe et al., 1969, "Inhibition of endogenous gastrin activity by incubation with antibodies to the C-terminal tetrapeptide of gastrin. Surgery 65: 5633-639 are different from the present invention in that the immunogen used was derived from the carboxyl terminal tetra-peptide amino acid sequence common to G₁₇, G₃₄, and to another important hormone, cholecystokinin ("CCK"). The immunogen of Jaffe et al. is thus of no practical value as an anti-gastrin vaccine component; on the contrary, it would produce a deleterious state in which all gastrin activity and other hormone function of G₁₇, G₃₄, together with CCK, would be blocked and eliminated by immunization.

This invention provides a novel immunological approach to the control and regulation of gastrin induced disorders such as peptic ulcers. According to the invention, antibodies are induced in the patient by active immunization with immunogens that selectively target specific forms of gastrin. Alternatively, the patient can be passively immunized with anti-gastrin antibodies specific for certain forms of gastrin.

In addition to peptic ulcers, other diseases appear to be related to the hormonal and stimulatory effects of gastrin. These diseases may also be treated by the selective anti-gastrin treatment of the invention.

An area of major medical importance for which the neutralization of gastrin hormonal activity has great therapeutic potential concerns the control of tumors and pathological conditions that are stimulated by gastrointestinal hormones. Several cancers of the gastrointestinal tract and associated tissues are stimulated to grow by the trophic action of gastrin. See Lamers, C.S.H.S., and Jansen, J.B.M.S., 1988, "Role of Gastrin and Cholecystokinin in Tumours of the Gastrointestinal Tract", Eur. J. Cancer Clin. Oncol. 22: 267-273. Gastrin promotes the growth of colon carcinoma, gastric carcinoma and gastric carcinoids. Gastrin antagonists may inhibit the growth of human colon cancer and enhance host survival as has been shown in mice; see, Beauchamp, R.O., et al., 1985, "Proglumide, A Gastrin Receptor Antagonist, Inhibits Growth Of Colon Cancer And Enhances Survival In Mice." Ann. Surg. 202: 303-309. The neutralization of gastrin tumor promoting activity may provide an important therapy for these diseases.

A second important application of gastrin neutralization therapy concerns conditions in which the hormone is overproduced. Certain cancers of the gastrointestinal tract, apudomas, produce extremely large quantities of gastrin. In either case, the excess hormone produced by the apudoma or pituitary tumor will have adverse physiologic effects on organs or tissues containing receptors for the hormone. Excess gastrin production by apudomas stimulates hypertrophy of the acid secreting epithelium of the stomach, leading to excess stomach acid secretion, peptic ulcer, and neoplastic changes in the epithelium.

Available treatment for tumors stimulated by gastrin and for tumors that produce gastrin consists primarily of surgical resection of the cancerous tissue. This approach is frequently unsuccessful; in many instances the tumors cannot be located or are present in anatomic sites that are inoperable. In most instances these tumors do not respond well to radiation or chemotherapy regimens. New treatments are needed to supplement present procedures.

A therapeutic method of selectively neutralizing the biological activity of these hormones would provide an effective means of controlling or preventing the pathologic changes resulting from excessive hormone production.

The method of cancer therapy described in this invention has several advantages over present treatment methods. The method is non-invasive, selectively reversible, does not damage normal tissue, does not require frequent repeated treatments, does not cross the blood brain barrier and has reduced side effects.

The therapy may be selectively reversed by injecting the patient with a pharmaceutical composition comprising a neutralizing epitope molecule. This molecule should comprise the epitope sequence free of an immunogenic carrier. This non-immunogenic molecule will bind to the free antibodies previously induced against the epitope in the host.

Brief Description of the Figures

Figure 1: illustrates the stomach acid secretions over time of a control rat injected sequentially with hG₁₇, antisera raised against an unrelated peptide and hG₁₇.

Figure 2: illustrates the stomach acid secretion over time of a rat injected sequentially with hG₁₇, hG₁₇ premixed with anti hG₁₇ antisera; and hG₁₇.

Figure 3: illustrates the stomach acid output over time in a rat actively immunized against G₁₇ in response to an injection of pentagastrin ("pG") followed by injections of G₁₇.

Figure 4: illustrates the stomach acid output over time in a rat actively immunized against G₁₇ in response to an injection of G₁₇ followed by injections of pentagastrin and G₁₇.

Figure 5: illustrates the stomach acid output over time in a control rat in response to the sequential injection of pG, G₁₇ and pG.

Figure 6: illustrates stomach acid output over time in a rat actively immunized against G₁₇ in response to sequential injections of G₃₄, G₁₇ and G₃₄.

Figure 7: illustrates stomach acid output over time in a control rat in response to sequential injections of G₃₄, G₁₇ and G₃₄.

Figure 8: depicts the binding capacity in picograms ("pg.") of Antigen per microliter ("ul") of sera of anti-G₁₇ antibodies induced by two synthetic peptide antigen epitope polymers of the invention.

Figure 9: illustrates the total quantity of stomach acid secreted by hG₁₇-immune rats and non-immune rats in response to graded doses of hG₁₇. Two non-immune rats and four hG₁₇-immune rats were injected with .12 micrograms ("ug") and 2.5 ug doses of hG₁₇. Two hG₁₇-immune rats were injected with doses of 25 ug and 250 ug of hG₁₇.

Figure 10: depicts the stimulation of the growth of human colon cancer cell line, HCT 116, by pentagastrin.

Figure 11: illustrates the effect of anti-G₁₇ antibodies on the growth of colon cancer implants in nude mice infused with G₁₇, as measured by mean tumor volume in cubic millimeters. Group I consisted of rats injected with rat anti-G₁₇ antibodies, and Group II consisted of rats injected with normal rat antibodies.

Figure 12: illustrates the effect of anti-G₁₇ on the growth of colon cancer implants in nude mice as measured by mean tumor volumes in cubic millimeters. Group 1 was injected with both G₁₇ and anti-G₁₇ antibodies. Group III was injected only with saline.

Description of the Invention

Since the different forms of gastrin vary in function, it is necessary to selectively neutralize specific forms of gastrin to control specific functions. To regulate gastrin mediated secretion of stomach acid following meals (the principal source of excess stomach acid relating to ulcers), an immunogen must specifically target G₁₇. In order to selectively neutralize G₁₇, one or more antigenic epitopes on G₁₇ that are not found on G₃₄ or cholecystokinin which exhibits carboxy terminal homology with gastrin must be identified. As discussed above even though the C-terminus of G₁₇ and G₃₄ are identical the N-terminus of G₁₇ is very different from that of G₃₄. This results in antigenic epitopes that are unique to G₁₇ and can be separately targeted. We have identified and mapped such a unique epitope on G₁₇. A specific embodiment of the present invention concerns immunogens comprising this unique epitope. These immunogens result in high levels of anti-G₁₇ antibodies that do not crossreact with G₃₄ and block some or all of G₁₇ stimulation of gastric acid secretion while still allowing G₃₄ and CCK, which share with G₁₇ a common receptor, to carry out their physiologic function. The regulation of acid secretions can also involve the neutralization of G₃₄; we have also identified and mapped unique epitopes on G₃₄ that are not found on G₁₇ or CCK.

Our immunoneutralizing approach has several attractive advantages over current treatments for peptic ulcer. One of these advantages is the overcoming of the major problem of patient compliance since a daily dose of a drug is not required. This invention treats ulcers by preventing the release of excess stomach acid, unlike antacids that neutralize secreted acid. By administering our synthetic peptide as an immunogen, the frequency and quantity of treatment administration is decreased, while at the same time long-lasting control of acid production, long term prevention of recurrence, and reduced side effects and easier patient administration are provided. Unlike conventional anti-ulcer drugs, antibodies generated by the peptide immunogens are very specific to their target. They do not cross the blood-brain barrier, and their use avoids certain complications encountered with drugs, for example, liver toxicities associated with H₂ antagonists. In addition, unlike this invention, agonists or antagonists of G₁₇ have reduced efficiency for controlling ulcers because such compounds have low specificity for the receptors for G₃₄ and CCK, which have identical receptor binding sites with G₁₇.

The immunogens against one form of gastrin, "little gastrin", or G₁₇, are constructed to produce an anti-gastrin immunogen component that will induce a selective and specific antibody response to G₁₇ in the immunized human or other vertebrate, but not to G₃₄ or CCK. This selective immunization to produce G₁₇ specific antibodies is crucial to avoid producing antibodies specific for or cross reactive with G₃₄, which might during the treatment of a specific condition induce undesirable side effects by blocking G₃₄ physiologic functions. The antibodies resulting from the immunization with such immunogens target the chemical structure of G₁₇ which is antigenically and immunogenically unique from the structure of G₃₄.

Peptides comprising the amino acid residues beginning from the amino terminus (amino acid residue number one) of G₁₇ and extending up to and including amino acid residue number 12 having the sequence pyro-Glu-Gly-Pro-Trp-Leu-Glu-Glu-Glu-Glu-Ala-Tyr, are used to prepare the immunogens of the invention by coupling them to an immunogenic carrier. For simplicity this sequence can be written based upon the international code for amino acids as pyro-Q-G-P-W-L-E-E-E-E-A-Y. The immunogens may contain a part or all of this sequence. The last 5 carboxy-terminal end amino acids of the G₁₇ chemical structure (residues 13-17) are preferably not used, because this sequence is a common antigenic sequence between G₁₇, G₃₄, and at least one other hormone, cholecystokinin (CCK). Fragments, extensions, or other subsets of the natural hormone and of this 12 amino acid sequence of G₁₇ may be used.

The most preferred peptide used in the invention is the hexamer pGlu-Gly-Pro-Trp-Leu-Glu-. The other peptides such as pGlu-Gly-Pro-Trp-Leu-Glu-Glu-Glu-, pGlu-Gly-Pro-Trp-Leu- and pGlu-Gly-Pro-Trp are also preferred. One or more other amino acids may also be substituted for dose of the natural sequence, so that increased or decreased binding capacity, specificity and/or titer of the antibody response against G₁₇ may be induced in the vaccinated host by the immunogen.

In other embodiments of the invention the use of preformed G₁₇ specific polyclonal and/or monoclonal antibodies and their derivatives or fragments produced by immunization, hybridoma, recombinant DNA or other technologies as a method of passive immunization for the control of gastric acid secretion stimulated by G₁₇ may be used.

The present invention also provides immunogens against a second form of gastrin, "big gastrin" or G₃₄. These immunogens are used to produce an anti-gastrin vaccine component that may be useful for the treatment or prevention of other gastrointestinal diseases and that will induce a selective and specific antibody response to G₃₄ (but not to G₁₇ or CCK) in the vaccinated human or other vertebrate. This selective immunization to produce G₃₄ specific antibodies is crucial to avoid producing antibodies specific for or cross reactive with G₁₇.

The G₃₄ immunogens specifically target chemical structures of G₃₄ which are antigenically and immunogenically unique from the structure of G₁₇. The chemical structures of G₃₄ utilized in this invention include, but are not limited to, peptides comprising the amino acid residues beginning from the amino terminus (amino acid residue number one) of G₃₄ and extending up to and including amino acid residue number 22. The sequence of this peptide is pyro-Glu-Leu-Gly-Pro-Gln-Gly-Pro-Pro-His-Leu-Val-Ala-Asp-Pro-Ser-Lys-Lys-Gln-Gly-Pro-Trp-Leu. Based upon the international code for amino acids, this sequence is pyroQ-L-G-P-Q-G-P-P-H-L-V-A-D-P-S-K-K-Q-G-P-W-L-. The G₃₄ immunogens may contain part or all of this sequence and comprise the sequence coupled to an immunogenic carrier. The sequence of the last 12 amino acids of the G₃₄ chemical structure (residues 23-34) are preferably not used in this invention because this sequence is a common antigenic sequence between G₁₇ and G₃₄. The sequence of amino acids are also not used since the sequence 29-34 has common antigenic sites with cholecystokinin. It is contemplated that the use of any fragments, extensions, or other subsets of the natural hormone and of this 22 amino acid sequence may be used as immunogens.

The most preferred peptide for use is the hexamer pGlu-Leu-Gly-Pro-Gln-Gly. The peptides pGlu-Leu-Gly-Pro-Gln-Gly-Pro-Pro-His- and pGlu-Leu-Gly-Pro-Gln-Gly-Arg- Pro-Pro-Pro-Cys are also preferred. One or more other amino acids may also be substituted and/or modified for those of the usual natural sequence, so that increased or decreased binding capacity, specificity and/or titer of the antibody response against G₃₄ may be induced in the vaccinated host by the immunogen.

Preformed G₃₄ specific monoclonal antibodies and their derivatives or fragments produced by hybridoma, recombinant DNA or other technologies may also be used as a method of passive immunization for the control of gastric acid secretion stimulated by G₃₄.

The peptides and immunogens may be produced by any process commonly used in the art including, for example, standard peptide synthesis technologies; methods employing recombinant DNA and associated technologies; antigen mimicking methods including antibody-internal image technology and any other related methodologies that produce a structure that immunologically resembles the antigenic structures of (mimotopes).

The means by which anti-gastrin antibodies prevent acid release has not been thoroughly established. Without being bound by theory, we believe that the acid suppressive effect of our immunogen is due to the binding of anti-gastrin antibodies to gastrin (G₁₇ and/or G₃₄) in the blood, and thereby preventing the binding of gastrin to its physiological receptors on the surfaces of parietal cells. Thus, gastrin is prevented from signaling parietal cells to secrete acid into the stomach.

For the G₁₇ and/or the G₃₄ epitopes of this invention to induce antibodies, it may be necessary to increase their immunogenicity by chemically coupling them to other molecules. Such molecules are termed "carriers". Any molecule capable of serving as a carrier may be used. Examples of carriers for this purpose

include: diphtheria toxoid, tetanus toxoid, keyhole limpet hemocyanin, bovine serum albumin, etc. Fragments of these carriers, including single epitopes, may also be used. Any method of chemically coupling the epitopes to the carriers may be followed. A preferred method utilizes the bifunctional linking agent ECMS described in U.S. Patent 4,302,386; Lee et al., 1981.

The epitopes can be alternatively rendered immunogenic by crosslinking (e.g., polymerizing) a number of epitopes. For this purpose, it may be necessary to extend the molecule bearing the G₁₇ or G₃₄ epitope by the addition of selected compounds that provide structures through which the crosslinking will occur. These additions must not disrupt the structure of the gastrin epitope, because the capacity to induce anti-gastrin antibodies would be lost. For example, to the carboxy terminal end of the G₁₇ epitope pyro-Q-G-P-W-M-E-E is added the amino acid sequence X-R-P-P-P-K, to give pyro-Q-G-P-W-M-E-E-K-R-P-P-P-K. This molecule is then polymerized with glutaraldehyde, which crosslinks the lysine K residues, to form the crosslinked immunogen. This crosslinked immunogen should induce specific antibodies against G₁₇.

It may be desirable in some applications to immunize against both G₁₇ and G₃₄. In this embodiment G₁₇ and G₃₄ immunogens are used in combination including optionally an immunogen with an epitope common to G₁₇ and G₃₄, e.g. Asp-Pro-Ser-Lys-Lys-Glu-Pro-Trp-Leu-, so that antibodies against both G₁₇ and G₃₄ are induced by the immunized host. The immunogens of this invention are therefore useful for more than just the treatment or prevention of ulcers. The immunogens may be used to treat any disease in which the gastrin stimulated secretion of stomach acid or stimulation of the growth of cancer by gastrin (e.g. colorectal and gastric) is a factor.

Administration of these immunogens, compositions containing them, or pharmaceutically acceptable and immunologically effective derivatives thereof, may be via any of the conventionally accepted modes of administration of agents which exhibit immunogenicity.

The compositions in which the immunogens are administered may be in a variety of forms. These include, for example, solid, semi-solid and liquid dosage forms, such as powders, liquid solutions or suspensions, suppositories, and injectable and infusible solutions. The preferred form depends on the intended mode of administration and therapeutic applications. The compositions also will preferably include conventional pharmaceutically acceptable carriers and may include other medicinal agents, carriers, adjuvants, excipients, etc., e.g., human serum albumin or plasma preparations. Preferably, the compositions of the invention are in the form of a unit dose. The amount of active compound administered as an immunization or as a medicament at one time, or over a period of time, will depend on the subject being treated, the manner and form of administration, and the judgment of the treating physician. However, an effective dose may be in the range of from about 1 µg to about 10 mg of the immunogen of this invention, preferably about 100 µg to about 2 mg; it being recognized that lower and higher doses may also be useful.

This invention provides a novel immunological approach to the treatment of tumors whose growth is dependent upon or stimulated by one or more of the forms of gastrin. According to the invention, antibodies are induced in the patient by passive or active immunization with immunogens that target one or more of the forms of gastrin. Such antibodies will bind to and neutralize the hormone to which the antibodies are directed, thereby preventing the hormone from acting upon the tumor. The antibody-mediated deprivation of hormonal activity thus constitutes a means of controlling gastrin dependent tumors.

This invention additionally provides a means of treatment for the effects of tumors that produce one or more forms of gastrin. In this embodiment, the immunogens of this invention are used to induce antibodies against the form of the hormone that is produced by the tumor. Such antibodies will bind to and neutralize the hormone, thereby preventing pathological consequences resulting from abnormally elevated levels of hormone.

This invention also provides a method for selectively reversing the antibody-mediated immunity induced by the anti-hormone immunogens of the invention. Soluble monovalent, monodeterminant epitopes are injected into the patient to bind to and neutralize the selected anti-hormone antibodies. The antibodies will then be incapable of binding additional quantities of hormone, and will no longer affect the hormone's biological activity.

The immunogens of this invention can be constructed to induce antibodies that specifically neutralize a single form of gastrin. This is advantageous in treating a tumor and pathologic condition, particularly one that is stimulated primarily or totally by a single form of gastrin. Alternatively, these specific immunogens can be selectively combined to produce an immunogen, in which the immunogen's individual constituents each induce antibodies that target distinct forms of gastrin. Combined immunogens would be used to treat tumors and conditions that are stimulated by more than one form of gastrin. Such an immunogen has the advantage of allowing for the addition or removal of individual immunogen constituents. The specificity of the antibody response can thus be controlled over time, thus enabling the physician to tailor the ongoing treatment to the needs of individual patients.

This example demonstrates a means of preparing immunogens to induce anti-G₁₇ or anti-G₃₄ antibody responses.

EXAMPLE 1

Peptides for the induction of specific immune responses to either G₁₇ or to G₃₄ were prepared by standard solid state synthesis methods. Each peptide was characterized as to amino acid content and purity.

Peptides with the following amino acid sequences were synthesized:

Peptide 1- Human G₁₇(1-6) ("hG₁₇(6)"): pGlu-Gly-Pro-Trp-Leu-Glu-Arg-Pro-Pro-Pro-Pro-Cys

Peptide 2- Human G₁₇(1-5) ("hG₁₇(5)"): pGlu-Gly-Pro-Trp-Leu-Arg-Pro-Pro-Pro-Pro-Cys

Peptide 3- Human G₁₇(1-4) ("hG₁₇(4)"): pGlu-Gly-Pro-Trp-Arg-Pro-Pro-Pro-Pro-Cys

Peptide 4- Rat G₁₇(1-6) Leu 5 ("rG₁₇(6)Leu 5"): pGlu-Arg-Pro-Pro-Leu-Glu-Arg-Pro-Pro-Pro-Pro-Cys

Peptide 5- Human G₃₄(1-6) ("hG₃₄(6)"): pGlu-Leu-Gly-Pro-Gln-Gly-Arg-Pro-Pro-Pro-Pro-Cys

Peptide 6- Human G₃₄(13-22) ("hG₃₄/G₁₇ combination"): Asp-Pro-Ser-Lys-Lys-Gln-Gly-Pro-Trp-Leu-Pro-Pro-Pro-Cys

Each of these peptides were conjugated to amino groups present on a carrier such as Diphtheria toxoid ("DT") via the terminal peptide cysteine residue utilizing heterobifunctional linking agents containing a succinimidyl ester at one end and maleimide at the other end of the linking agent.

To accomplish the linkage between any of Peptides 1-6 above and the carrier, the dry peptide was dissolved in 0.1M Sodium Phosphate Buffer, pH 8.0, with a thirty molar excess of dithiothreitol ("DTT"). The solution was stirred under a water saturated nitrogen gas atmosphere for four hours. The peptide containing reduced cysteine was separated from the other components by chromatography over a G10 Sephadex column equilibrated with 0.2M Acetic acid. The peptide was lyophilized and stored under vacuum until used. The carrier was activated by treatment with the heterobifunctional linking agent e.g. Epsilon-maleimidocaproic acid N-hydroxysuccinimide ester, ("EMCS"), in proportions sufficient to achieve activation of approximately 25 free amino groups per 10⁵ molecular weight of carrier. In the specific instance of diphtheria toxoid, this amounted to the addition of 6.18 mg of EMCS (purity 75%) to each 20 mg of diphtheria toxoid.

Activation of diphtheria toxoid was accomplished by dissolving each 20 mg aliquot of diphtheria toxoid in 1 ml of 0.2 M Sodium Phosphate Buffer, pH 6.45. Aliquots of 6.18 mg EMCS were dissolved into 0.2 ml of Dimethyl Formamide ("DMF"). Under darkened conditions, the EMCS was added dropwise in 50 microliter ("ul") amounts to the DT with stirring. After 2 hours of incubation in darkness, the mixture was chromatographed on a G50 Sephadex column equilibrated with 0.1 M Sodium Citrate buffer, pH 6.0, containing 0.1 mM EDTA.

Fractions containing the EMCS activated diphtheria toxoid were concentrated over a PM 10 ultrafiltration membrane under conditions of darkness. The protein content of the concentrate was determined by either the Lowry or Bradford methods. The EMCS content of the carrier was determined by incubation of the activated carrier with cysteine-HCl followed by reaction with 10 mM of Elman's Reagent 5,5 dithio-bis (2-nitrobenzoic acid) 10mM. The optical density difference between a blank tube containing cysteine-HCl and the sample tube containing cysteine-HCl and carrier was translated into EMCS group content by using the molar extinction coefficient of 13.6×10^3 for 5-thio-2-nitro benzoic acid at 412 nm.

The reduced cysteine content (-SH) of the peptide was also determined utilizing Elman's Reagent. Approximately 1 mg of peptide was dissolved in 1 ml of nitrogen gas-saturated water and a 0.1 ml aliquot of this solution was reacted with Elman's Reagent. Utilizing the molar extinction coefficient of 5-thio-2-nitro-benzoic acid (13.6×10^3), the free cysteine -SH was calculated. An amount of peptide containing sufficient free -SH to react with each of the 25 EMCS activated amino groups on the carrier was dissolved in 0.1M Sodium Citrate Buffer, pH 6.0, containing 0.1 mM EDTA, and added dropwise to the EMCS activated carrier under darkened conditions. After all the peptide solution had been added to the carrier, the mixture was incubated overnight in the dark under a water saturated nitrogen gas atmosphere.

The conjugate of the peptide linked to the carrier via EMCS is separated from other components of the mixture by chromatography over a G50 Sephadex column equilibrated with 0.2 M Ammonium Bicarbonate. The conjugate eluted in the column void volume is lyophilized and stored desiccated at -20° C until used.

The conjugate may be characterized as to peptide content by a number of methods known to those skilled in the art including weight gain, amino acid analysis, etc. Conjugates of Peptides 1-6 and diphtheria toxoid produced by these methods were determined to have 20-25 moles of peptide per 10⁵ MW of carrier

and all were considered suitable as immunogens for immunization of test animals.

EXAMPLE 2

As examples of the utilization of peptides containing sequences of human gastrin as immunogens to induce immune responses against hG₁₇ or hG₃₄, we have immunized rats with the conjugate immunogens constructed from Peptides 1-6 of Example 1 and Diphtheria toxoid ("DT") (referred to as Immunogens 1-6, respectively).

Six different groups of 15 Sprague-Dawley female rats (200 gm. body weight) were each immunized with one of the immunogens constructed from Peptides 1-6. Each animal was injected subcutaneously with 0.25 ml of immunogen consisting of 0.1 mg of conjugate dissolved in 0.125 ml of 0.1 M Sodium Phosphate Buffered Saline, pH 7.3, emulsified with an equal volume of Squalene-Arlacel (4:1 ratio volume/volume) vehicle containing 0.05 mg of Nor MDP as adjuvant.

Two additional groups of 15 rats were immunized with a peptide-DT conjugate in which the peptide had no sequence homology with gastrins so as to act as a negative immunization control.

Each rat was given an injection of immunogen at 0, 3, and 6 weeks. Blood was collected from each rat at 3, 6, and 8 weeks of the experiment. Serum was collected from each blood sample and stored at -20° C until utilized in assays to determine the presence of anti-gastrin antibodies.

Two types of assays were used to detect anti-gastrin antibodies. A solid-phase enzyme linked immunosorbent assay (ELISA) and a liquid phase radioimmunoassay (RIA) were employed.

ELISA was used to screen for reaction or cross reaction of antisera raised against Peptides 1-6 with Peptides 1-6 or with hG₁₇, hG₃₄, or hCCK. The RIA was used to quantitate the antibody levels in the antiserum of each immunized animal that was reacted with hG₁₇ or hG₃₄ by determining the antigen binding capacity (ABC), expressed as pg hormone bound per ("ul") of antiserum (pg/ul).

The ELISA was conducted by coating polystyrene 96 well plates (Immulon II) with 1 ug/ml of Peptides 1-6, hG₁₇, hG₃₄, or hCCK antigen. Serial dilutions of test antisera of 1×10^{-4} to 1×10^{-8} were incubated with each test peptide for 30 minutes at room temperature. In some instances antisera raised against a particular peptide of the Peptides 1-6 were preincubated with large excesses of the other peptides of the Peptide 1-6 group or with hG₁₇ or hG₃₄ in an attempt to inhibit binding of the antiserum to its particular peptide and also to demonstrate the occurrence of antibodies in the antisera that were specific for the sequence (spacer) of each peptide that was common to all of peptides 1-6 (e.g. Arg-Pro-Pro-Pro-Cys). After washing each well thoroughly to remove unbound antibody, each well was treated with biotinylated anti-rat immunoglobulin reagent for 30 minutes at room temperature. After another wash sequence to remove unbound anti-rat reagent, avidin-alkaline phosphatase conjugate was added and the mixture was incubated for an additional 30 minutes. The mixture was washed thoroughly to remove unbound avidin-alkaline phosphatase reagent, and the chromogenic substrate PNPP was added for a 10 minute period. The absorbance of each well was read at 490 nm after the 10 minute incubation.

The standard RIA procedure was followed. In the RIA, 0.1, 1.0 or 10.0 ul aliquots of antiserum were incubated with approximately 200 pg of ¹²⁵I labeled hG₁₇ or 400 pg of labeled hG₃₄. The antisera were incubated with label for 2 hours, and were followed by a precipitation of hormone-antibody complexes with 25% polyethylene glycol. Antigen binding capacities for each antiserum were then determined from the amount of radioactive hormone precipitated. To demonstrate the specificity of the reaction of the ¹²⁵I labeled hormone with the antisera, aliquots of the antisera were preincubated in some tests with excess amounts of the hormone that were not labeled with ¹²⁵I to inhibit binding of the antisera to the labeled hormone.

The specificities of the antibody responses induced by Immunogens 1-6 as measured by ELISA are depicted in Table 1. Immunogen 1, containing the peptide sequence of hG₁₇ (1-6), induced antibodies that reacted strongly with hG₁₇ and hG₁₇ (1-6) peptide, but only weakly with hG₃₄ (1-6) or hG₃₄ (13-22). Antisera raised to Immunogen 1 did not react with hG₃₄. Inhibition experiments with Peptides 1-6 demonstrated that the weak reactivity of anti-Immunogen 1 antibodies with hG₃₄ (1-6) and G₃₄ (13-22) peptides was due to the presence of antibodies that were induced by the spacer sequence (-Arg-Pro-Pro-Cys) common to all the peptide sequences of Immunogens 1-6.

Immunogens 2 and 3 induced antibody responses specific for hG₁₇ that were much weaker than those induced by Immunogen 1 (Table 1). Inhibition experiments, demonstrated that the weak reactivities of anti-Immunogen 2 and 3 antibodies for Peptides 1-6 are specific for the common spacer sequence of Peptides 1-6.

Immunogen 4, containing the rat G₁₇ sequence, induced antibodies that weakly reacted with Peptides 1-6, but not hG₁₇ or hG₃₄ (Table 1). Inhibition experiments demonstrated that these antibodies were directed against the spacer sequence common to Peptide 1-6.

Immunogen 5 hG₃₄ (1-6) , induced antibodies that strongly reacted with hG₃₄ and Peptide 5, hG₃₄ (1-6) , but weakly with the other peptides and not at all with G₁₇. Inhibition experiments demonstrated that the reactivity with peptides 1-4 and 6 was, due to anti-spacer specific antibodies.

Immunogen 6, hG₃₄ (13-22), induced antibodies that reacted weakly with Peptides 1-6, but not with hG₁₇. Inhibition experiments demonstrated that the antibodies binding Peptides 1-6 were specific for the common spacer sequence.

10

15

20

25

30

35

40

45

50

55

Table 1

Antisera to:		Reaction in ELISA with:						
	hG ₁₇	Peptide 1 hG ₁₇ (1-6)	Peptide 2 hG ₁₇ (1-5)	Peptide 3 hG ₁₇ (1-4)	Peptide 4 rG ₁₇ (1-6)	Peptide 5 hG ₃₄ (1-6)	Peptide 6 hG ₃₄ (13-22)	HG ₃₄
Immunogen 1 hG ₁₇ (1-6)-DT	++	+++	+	±	±	±	±	0
Immunogen 2 hG ₁₇ (1-5)-DT	+	±	+	±	±	±	±	0
Immunogen 3 hG ₁₇ (1-4)-DT	±	±	±	+	±	±	±	0
Immunogen 4 rG ₁₇ (1-6)-DT	0	±	±	±	+	±	±	0
Immunogen 5 hG ₃₄ (1-6)-DT	0	±	±	±	±	++	±	+++
Immunogen 6 hG ₃₄ (13-22)-DT	±	±	±	±	±	±	+	±
+++ to ++: Strongly reactive								
±: Weekly reactive								
0: No reaction								

All antisera were also tested against hCCK; none of the antisera bound to hCCK.

Table 2 demonstrates the RIA-measured antigen binding capacities ("ABC") versus hG₁₇ or hG₃₄ of antisera raised against Immunogens 1-6 after three immunizations of rats with 0.1 mg of conjugate.

Table 2

Rats Immunized With:	Mean RIA ABC (pg/ul)	
	hG ₁₇	hG ₃₄
Immunogen 1 hG ₁₇ (1-6)-DT	19.29	0.00
Immunogen 2 hG ₁₇ (1-5)-DT	7.59	0.00
Immunogen 3 hG ₁₇ (1-4)-DT	2.15	0.00
Immunogen 4 rG ₁₇ (1-6)-DT	0.00	0.00
Immunogen 5 hG ₃₄ (1-6)-DT	0.00	6.38
Immunogen 6 hG ₃₄ (13-22)	0.00	1.28

The liquid phase RIA demonstrated that Immunogens 1-3 containing the hG₁₇ peptide sequence induced antibodies that reacted only with hG₁₇ and that Immunogen 5 containing the hG₃₄ sequence, induced antibodies that reacted only with G₃₄. Immunogen 6 induced very low ABC's to G₃₄.

The ELISA and RIA assays thus demonstrate the specificity of the responses to hG₁₇ or hG₃₄ that are induced by Immunogens 1-6.

EXAMPLE 3

This example demonstrates the ability of antisera raised against Peptide 1 (hG₁₇ (1-6)) to neutralize the in vivo acid stimulating activity of hG₁₇. In this demonstration an amount of hG₁₇ is mixed with an excess amount of anti-Peptide 1 antiserum sufficient to bind to all the hG₁₇ prior to injection of the complex into a normal (non-immunized) rat.

In control experiments the amount of hG₁₇ sufficient to stimulate an increase of acid secretion of at least 100% above nonstimulated acid secretion in normal rats was determined to be 0.4 ug of hG₁₇ hormone per kg body weight.

Antisera from the rats immunized with Immunogen 1 were pooled and standard amounts of antisera were incubated with 200 pg ¹²⁵I labeled hG₁₇ after incubation with increasing amounts of cold hG₁₇ as inhibitor. Based on this inhibition study 1 ml of antiserum was capable of binding 1000X the 0.4 ug/kg dose of hG₁₇ to be administered to rats. As a safety factor, the 0.4 ug/kg (approximately 120 ng) of hormone was mixed with 2.5 ml of anti-hG₁₇ specific antiserum raised against Immunogen 1.

Rats to be injected with hG₁₇ complexed with anti-hG₁₇ antibodies were surgically prepared for collection of stomach secretions by the perfused rat stomach procedure.

Under general anesthesia and following tracheostomy, the rat was cannulated via the esophagus and duodenum to allow continuous perfusion of the stomach with 0.9% saline. The stomach perfusate was collected as 5 minute interval samples and was titrated for acid content by neutralization with base (sodium hydroxide). Incremental and total acid input during the duration of the experiment and after each treatment was determined.

Each control or experimental test rat was first injected with 0.4 ug/kg hG₁₇ to determine the rats total acid secretory response to this treatment. The first treatment was followed one hour later in test rats with an injection of 0.4 ug/kg of hG₁₇ that had been premixed for one hour with 2.5 ml of anti-hG₁₇ specific antiserum. Control rats received an injection of hG₁₇ mixed with 2.5 ml of antiserum raised against an unrelated peptide. After one hour, a second injection of free hG₁₇ was administered to the test and control rats; and stomach perfusate was collected for an additional hour. The total acid output induced by the second and third injections of hG₁₇ were expressed as a percentage of the total acid output induced by the first injection of hG₁₇.

In five rats tested by this experimental procedure here was an 81%-100% (mean = 94%) reduction in

the acid secreted by the perfused rat stomach in response to the hormone premixed with anti-hG₁₇ specific antibody (second injection) or to the third injection consisting of free hG₁₇ alone. Control rats experienced little or no reduction in acid secretion stimulated by the second and third injections of hormone. Figure 1 and Figure 2 illustrate the responses of a control rat (Figure 1) and experimental rat (Figure 2) to these treatments.

EXAMPLE 4

A major application of this invention is the active immunization of humans to induce specific immunity against G₁₇ for ulcer therapy and prevention. In this example, it is demonstrated that active immunization with an anti-G₁₇ immunogen induces antibodies that dramatically suppress G₁₇ mediated release of stomach acid.

To actively immunize rats against G₁₇, we follow the methods used to obtain antisera in the passive immunization tests as described in Example 3. An immunogen consisting of the G₁₇(6) peptide covalently coupled to Diphtheria toxoid (DT) is prepared as described in Example 1. This immunogen is suspended in Phosphate Buffered Saline at a concentration of 4.0 mg/ml. The antigen is emulsified in squalene:arlacel (4:1) vehicle, at a final ratio of 1:1 (antigen:vehicle). Nor-MDP is included in the mixture to give a final concentration of nor-MDP of 200 ug/ml. The final concentration of the DT-G₁₇(6) in the formulation is 2.0 mg/ml. Experimental rats are injected with 0.25 ml of this preparation intraperitoneally. Each injection thus delivers approximately 500 ug of immunogen plus 50 ug of nor-MDP. A second injection is similarly administered 21 days later.

Blood samples for antibody analysis are obtained by tail vein bleeding before the first injection and 14 days after each injection. Sera is prepared by allowing the blood to clot for 30 minutes at room temperature followed by centrifugation at 400 x g to remove the clots. The Sera are stored frozen until used.

To determine the antibody responses of the immunized rats, a RIA is employed as described in Example 2. The results of this test show that the immunization procedure induces high titers of antibody against G₁₇. These responses are specific for G₁₇; no reactivity is detected with G₃₄, with pentagastrin (the biologically active, carboxy terminal fragment of G₁₇, G₃₄, and CCK), or with CCK. The antibodies are thus directed against the unique epitope on G₁₇ that is selectively targeted by the immunogen. These results are similar to those of Example 2.

The use of the immunogens described herein for the active immunogen is not limited to the adjuvant, vehicle, injection schedule, etc., described above. Any means of safely inducing immunity against G₁₇ using the immunogens described can be applied. This includes using alternative dosages, routes, vehicles, adjuvants, excipients, slow-release compounds, etc.

We test for the neutralization of G₁₇'s biological activity in the immunized animals using the perfused rat stomach method, as described in Example 3, with the important difference that we do not inject antisera into the rats (passive immunization) because the actively immunized rats are making their own antibodies against G₁₇. The dosages of compounds administered in these tests, with delivery times of 5 minutes per total dose, are: G₁₇ = 0.4 ug/kg, G₃₄ = 0.8 ug/kg, and pentagastrin = 2.0 ug/kg. Stomach contents sampling times are 5 minutes per sample. The stomach acid outputs are calculated as the percent of maximal acid output

$$= (100) \frac{A_n - A_b}{A_{max} - A_b},$$

where A_n = the acid produced over each 5 minute sampling interval (as determined by titration with NaOH); A_{max} = the maximal 5 minute release of stomach acid upon stimulation, usually (but not necessarily) by pentagastrin; and A_b = the baseline level of acid present at the time of a given stimulation (with G₁₇, pentagastrin, or G₃₄).

The effects of active immunization against G₁₇ upon the G₁₇ and pentagastrin ("pG") induced acid secretion are shown in Figures 3 and 4. The ordinate represents the percent of acid output compared to the maximal acid output induced by pentagastrin. These experiments differed, by design, in the order of G₁₇ and pentagastrin challenge. In both cases, it is clear that in the G₁₇ immunized rats the production of stomach acid in response to G₁₇ (Figure 3, Peaks 2 and 3; Figure 4, Peaks 1 and 3) is substantially reduced in comparison with acid secretion induced by pentagastrin (Figure 3, Peak 1; Figure 4, Peak 2).

The mean reduction in the total G_{17} mediated acid secretion in our G_{17} immune rats is 85% (compared to pentagastrin).

We verified that the acid reductions were a direct consequence of immunization against G_{17} by conducting challenges with G_{17} or pentagastrin in control rats. The control animals were immunized in an identical manner as the G_{17} -immune rats, except that the controls received antigen consisting of DT conjugated to an unrelated peptide (i.e., non-crossreactive with gastrin). RIAs and ELISAs, run on sera from these animals, demonstrated that they produced high antibody titers against both DT and the unrelated peptide, but none against G_{17} , pentagastrin, G_{34} or CCK. When tested for acid secretion, the control rats responded equally well to challenges with both G_{17} and pentagastrin. The results of such a test are shown in Figure 5. This rules out the remote possibility that the neutralization of G_{17} in the G_{17} -immune rats was caused by non-specific factors (e.g., adjuvant effects, crossreactive anti-DT antibodies, etc.).

A technical challenge presented by the perfused rat stomach assay was the selection of the appropriate acid stimulatory compound for use as a positive control. The exquisite specificity for G_{17} 's unique epitope that is characteristic of antibodies induced by our immunogen enabled us to use the ideal control compound: pentagastrin. Pentagastrin comprises the receptor binding/stimulatory sequence of G_{17} and also of both G_{34} and CCK, and it is not bound by antibodies induced by our immunogen. The responses to pentagastrin demonstrated that our immunized animals' acid response mechanism to G_{17} stimulation were functional. In addition, the pentagastrin responses established the level of acid secretion to be expected from G_{17} stimulation. The dosages of G_{17} and pentagastrin, which we determined experimentally, were selected to induce approximately equal acid secretory responses in control rats (see Figure 5). Thus, we were able to accurately quantitate reductions in acid secretion resulting from the neutralization of G_{17} .

For completeness, we have also challenged with G_{34} . We designed our immunogen to specifically neutralize G_{17} mediated acid secretion (particularly following food intake) and to have no effect upon acid output induced by G_{34} (which provides for basal stomach activity). Since the antisera from the G_{17} immune rats do not react with G_{34} , we expected to see no effect upon G_{34} 's ability to stimulate acid secretion. Indeed, as shown in Figure 6, the immunized rats secreted normal quantities of acid in response to G_{34} stimulation (Peaks 1 and 3). As expected, the injection of G_{17} failed to induce acid secretion in these animals (Figure 6, Peak 2). Both G_{17} and G_{34} induce strong acid secretory responses in control rats (immunized against an irrelevant peptide), as can be seen in Figure 7. Clearly, the anti- G_{17} antibodies induced by our immunogen have no effect upon the functions of other molecules to which the antibodies do not bind. The $G_{17}(6)$ based immunogen described herein induces antibodies that are specific for G_{17} and neutralize G_{17} 's acid releasing activity. Such an immunogen should thus protect against and cure peptic ulcers.

EXAMPLE 5

This example demonstrated that a polymerised peptide immunogen can be constructed and used to reduce anti- G_{17} antibody responses. Synthetic peptides have been produced that contain the unique epitope on G_{17} and in addition carry reactive groups that can be selectively bound to crosslinking agents. These peptides serve as monomers in the construction of a polymer immunogen. By including two or more reactive groups in each peptide it is possible to construct multi-peptide aggregates, or polymers, by reaction of the groups with a cross-linking agent. Such polymers are then used as immunogens to induce antibodies against the G_{17} epitope expressed by the peptide. These antibodies bind to G_{17} *in vivo* and neutralize G_{17} , thus mediating an anti-ulcer effect. These polymerized peptides have an advantage in that they can be used as immunogens by themselves without a coupling to an immunogenic carrier.

The following peptide designated as Peptide 7 was constructed:

1 2 3 4 5 6 7 8 9 10 11 12 13 14
pGlu-Gly-Pro-Trp-Leu-Glu-Glu-Lys-Arg-Pro-Pro-Pro-Pro-Lys

The G_{17} epitope is contained in amino acids 1-7 of the peptide. Other epitopes eg. G_{34} epitopes, can also be used to construct other polymer immunogens according to the invention. Amino acids 8 and 14, which are both Lys, contain amino groups as side groups. These amino groups act as functional groups which are reacted with the functional groups on the crosslinking agent to form the crosslinked peptide polymer. Other amino acids containing side functional groups could be substituted for Lys depending on the

reactivity of the functional group with the group on the crosslinking agent to be used. The location of the functional amino acids can be varied in the peptide as long as they are not positioned within the epitope region. Additional reactive amino acids could also be added to increase crosslinking. These additional amino acids could be reactive with the same or alternative crosslinking agents. It follows that more than one type of crosslinking agent can be used.

Amino acids 9-13 comprise a "spacer region" between the reactive amino acids 8 and 14. The composition, number of amino acids and length of the spacer can be varied. If desirable, helper T-cell epitopes can also be included in the peptide.

Peptide 8 was synthesized and purified by standard solid phase peptide synthesis and purification methodologies. Any other method of peptide production well known to those skilled in the art including recombinant DNA technology can also be used to produce the peptides of the invention.

5.0 mg of the peptide was dissolved in 1.0 ml phosphate buffer (0.1M; pH=6.8). To this was added glutaraldehyde (Grade 1, Sigma Chemical Co.) in a 2:1 molar ratio of glutaraldehyde to peptide. The glutaraldehyde was added dropwise with stirring, at room temperature.

The reaction was allowed to proceed overnight, at room temperature, with stirring. 50.0 mgs. of sodium borohydride were then added slowly to the reaction mixture, and the mixture was stirred at room temperature for an additional hour. The mixture was transferred to dialysis tubing, 1,000 molecular weight cutoff (#132636, Spectrum Medical Industries, Inc.), and exhaustively dialyzed against saline. The peptide-polymer was stored frozen at -20°C.

The polymer was analyzed by SDS-PAGE using a 15% polyacrylamide gel. The electrophoresis demonstrated that the polymerized peptide contained polymers of various sizes comprising multiples of the peptide. The average polymer contained 6 peptides, however; the size of the polymers ranged up to 12 peptides per molecule.

A second polymer was made using the identical procedures, except that a 20:1 molar ratio of glutaraldehyde to peptide was used. The SDS-PAGE analysis of the second polymer similar results with respect to the size range as compared to the 2:1 polymer.

Each polymer preparation, 2:1 and 20:1, was used to immunize two separate groups of five mice per group. prior to injecting the mice with polymer, blood samples were taken from each mouse. The preparation was suspended in Freund's Complete Adjuvant ("FCA") H37Ra (DIFCO Labs) in a 1:1 (vol:vol) ratio of polymer:FCA. The mice were each injected intraperitoneally with 100 ug polymer in 0.2 ml of the mixture. After 21 days each mouse was given a second injection of the same polymer with which it had been injected previously. In the second injection, the antigen was administered intraperitoneally in saline, at 100 ug per mouse. Each mouse was bled 14 days after the second injection and the sera were isolated. The mouse sera were assayed for anti-G₁₇ antibodies by radioimmunoassay (RIA). 1.0 ul of sera was added to 300 ul of buffer (1% BSA in phosphate buffered saline with 0.005M EDTA, pH=7.2). To each of these samples was added 100 ul or 3000 CPM of ¹²⁵I-labeled G₁₇ (NEN, Specific activity = 12 uCi/ug). The samples were incubated 1.0 hour at room temperature. We next added 100 ul of Calf Serum (Hyclone Labs), immediately followed by 500 ul of 25% polyethylene glycol-8000 (Sigma). The samples were mixed and then centrifuged for 30 minutes at 500 x g at room temperature. The supernatant was discarded, and the pellet suspended in 250 ul of saline at 90°C. The suspension was transferred to 3.0 ml of Scintiverse II [Fisher Scientific] in mini vials for liquid scintillation counting. The samples were counted in a Beckman Liquid Scintillation counter (#LS 5000 LE) for ¹²⁵I. The binding capacities of the antisera were calculated from the resulting ¹²⁵I counts per sample and are depicted in Figure 8.

Both of the polymers induced anti-G₁₇ antibody responses. Polymer #1, the 2:1 ratio polymer (Figure 8), induced a very strong response of 56 pg of antigen bound per ul of sera. Polymer #2, the 20:1 ratio polymer (Figure 8), induced a response that was 10-fold lower. The response induced by polymer #1 is equivalent to that induced by three injections of the G₁₇(6)-DT immunogen of Example 2 in rats.

Thus, polymerized synthetic peptides can be used to induce potent anti-G₁₇ antibody responses.

EXAMPLE 6

The following example demonstrates that anti-G₁₇ antibodies neutralize the tumor stimulatory activity of G₁₇ in vitro.

HCT-116 cells (a human colon cancer cell line) were cultured in McCoy's 5a medium (McCoy et al., Proc. Soc. Exper. Biol. Med. 100:115-118) supplemented with epidermal growth factor (10 ng/ml), insulin (20 ug/ml), transferrin (4 ug/ml), sodium selenite (10⁻⁸M), hydrocortisone (2 ug/ml), and triiodothyronine

(4×10^{-10} M). Subcultures were made once weekly for four weeks by treating cultures with 0.5% Trypsin + 0.2% EDTA in Hanks Balanced Salts Solution to remove adherent cells, followed by inoculation of T-75 tissue culture flasks with approximately 1×10^6 cells. Cultures were maintained under standard conditions (37° , 100% humidity, 5% CO_2).

5 Prior to testing, the HCT-116 cells were synchronized to late G_0 phase with thymidine, as follows: the HCT-116 cells were seeded into 24 well culture plates at approximately 1×10^4 cells per well and incubated overnight in 1 ml supplemented McCoy's medium 5a. The medium was then removed and replaced with fresh supplemented medium. Thymidine was added to 0.8 mM final concentration to each well and the cultures were incubated for 24 hours. At the end of the synchronization period, the medium containing
10 thymidine was replaced with test media as described below.

To demonstrate that HCT-116 cells proliferate under the influence of gastrin, the synchronized HCT-116 cells were grown in supplemented McCoy's 5a medium, in the presence or absence of 10 μM pentagastrin (the hormonally active segment of gastrin). To assess cell proliferation, total cell counts were performed after selected incubation times. As shown in Figure 10, HCT-116 cells proliferated more rapidly in the
15 presence of pentagastrin than in the absence of the hormone. This difference was already evident after three days of culture, and was maximal by day five. The increase in the number of cells of pentagastrin treated culture at the end of eight days was three times greater than that of the non-pentagastrin treated cultures.

To demonstrate that anti- G_{17} antibodies neutralize the proliferative activity of G_{17} , the effect of anti- G_{17} antisera upon G_{17} -induced increases in the rate of [^3H]-thymidine uptake by HCT-116 cells was studied.
20

Synchronized HCT-116 were cultured in 24 well plates of 10^4 cells/well in supplemented McCoy's 5a medium without FBS. The total culture volume was 1 ml per well. Four wells were cultured for each test condition. Cells were grown in presence of G_{17} at two concentrations, 5 μM and 50 μM . At each G_{17} concentration, rat anti-human G_{17} antisera (antigen binding capacity determined by RIA = 30 $\mu\text{g}/\mu\text{l}$) or
25 normal rat sera were added at a final dilution of 1:25.

After 8 days of culture, 0.4 μCi of [^3H] thymidine (specific activity = 2 Ci/mMole) were added to each test well. Following 16 hours incubation, the replicates were processed with a Multiple Automated Cell Harvester (Mini-Mash II, Whittaker Bioproducts) and the [^3H]-thymidine incorporation was determined by scintillation counting.

30 As shown in Table 3, the addition of anti- G_{17} antiserum caused a substantial reduction in [^3H]-thymidine incorporation, relative to uptake by cells cultured in the presence of normal rat serum. At 5 μM G_{17} , counts incorporated were reduced by 59%; at 50 μM G_{17} , label uptake was reduced by 34%. As [^3H]-thymidine incorporation directly reflects cell proliferation, this test shows that anti- G_{17} antiserum inhibits the proliferation activity of human G_{17} on colon cancer cells.

35 Table 3

Quantity of Human G_{17} in Culture	Mean CPM [^3H] Thymidine Incorporated(\pm SE)	
	Normal Rat Serum	Anti-h G_{17} Serum
50 μM	2130 \pm 141	1404 \pm 310
5 μM	1621 \pm 131	664 \pm 206

50 EXAMPLE 7

The following experiment was performed to demonstrate that the growth of established HCT-116 tumors is retarded when nude mice bearing the tumors are treated with anti-gastrin immunoglobulin.
55

Anti-human G_{17} serum was obtained from rats immunized against h $G_{17}(6)$ -DT (i.e., Immunogen 1). Normal serum was obtained from nonimmunized rats. Immunoglobulin fractions of the anti- G_{17} serum and the normal rat serum are prepared by affinity chromatography, using Protein A Sepharose. Both im-

munoglobulin preparations were adjusted to a final protein concentration of 1 mg/ml in PBS. Measured by RIA, the G₁₇-direct antigen binding capacity (ABC) of the anti-human G₁₇ immunoglobulin preparation was 30 pg/ul. The normal rat immunoglobulin preparation had no anti-human G₁₇ activity.

Thirty nude mice were each implanted subcutaneously (dorsally near the left shoulder) by trocar needle (14 gauge) with a single 2 mm cube of HCT-116 tumor tissue. The tumors were allowed to grow for one week prior to random assignment of individual mice to one of three treatment groups, ten mice per group, on day 0 of the Test.

Group I was treated with anti-human G₁₇ immunoglobulin function. Each mouse in Group 1 received intraperitoneal injections of 0.5 mg. of immunoglobulin beginning on day 0 and repeated on days 4, 8, and 14 of the test. The Mean human G₁₇ ABC of the mouse sera on day 16 was 8.3 ± 1.3 pg/ul.

Group II was treated with normal rat immunoglobulin fraction. Each mouse received intraperitoneal injections of 0.5 mg of immunoglobulin beginning on day 0 and repeated on days 4, 9, and 14, of the test. On day 16, the human gastrin G₁₇ ABC's of those sera was opg/ul.

Group III received no immunoglobulin injection but was injected instead with saline on days 0, 4, 9, and 14 of the test. The human gastrin G₁₇ ABC's of those sera was Opg./ul. on day 16.

On day 1, mice of Groups I and II were implanted subcutaneously with osmotic pumps (Alzet 2002) that were charged with human G₁₇ that delivered 10 ug/day human G₁₇ continuously for 14 days. On day 1, mice of Group III were implanted subcutaneously with osmotic pumps charged with saline which delivered a dose of saline instead of hormone continuously for 14 days.

Each mouse was observed daily for changes in the size of their individual tumors. Measurements of the tumors were made by vernier caliper approximately every other day. Volumes of tumors were estimated by the calculation: volume = (length X width²) ÷ 2 (Euhus et. al., 1986, J. Surg. Oncol. 31:229-234).

Observations on Group II mice were made through day 17 of the test, on which day the Group II mice were euthanized. Observations on Groups I and III were continued until day 32 of the test.

As shown in Figure 11, tumors in Group II mice (administered human G₁₇ and treated with normal rat immunoglobulin) grew very rapidly in response to the added gastrin, increasing more than 120 fold in volume in 16 days.

Tumors of Group I mice (administrated human G₁₇ and treated with anti-human G₁₇ immunoglobulin) grew at a significantly slower rate than the tumors of Group II. By day 16, the volume of the tumors of Group I mice were, on average, approximately 11 times smaller than those of Group II mice (Table 4). The results of this test demonstrate that HCT-116 tumors are stimulated to grow by human gastrin G₁₇ and that treatment with anti-human G₁₇ specific immunoglobulin neutralizes this growth-promoting effect and significantly slows the growth of HCT-116 tumors.

Tumors of Group III mice (not administered hG₁₇ and no immunoglobulin treatment) grew at a faster rate than Group I (Figure 12), suggesting an autocrine production of human G₁₇ (by the HCT-116 tumor cells) that stimulated the tumors to grow. Tumor-produced G₁₇ would be neutralized in the Group I animals, due to the injected anti-G₁₇ immunoglobulins. On day 32 of the test, the Group III tumors had attained approximately the same volume attained by Group II tumors on day 16. However, on day 32 tumors of Group I were significantly smaller than tumors of Group III (i.e., 3.3 times smaller volume, Table 4), indicating an inhibitory effect of the anti-G₁₇ immunoglobulin on tumor growth.

Table 4

Group	Treatment	Mean Tumor Volume (mm ³ ± SE)	
		day 16	day 32
I	Anti-G ₁₇ Antibody plus G ₁₇	142 ± 38	545 ± 137
II	Normal Rat Serum plus G ₁₇	1512 ± 348	-
III	Saline and no G ₁₇	391 ± 63	1825 ± 313

EXAMPLE 8

The following test demonstrates that antibodies against human G₁₇ inhibit tumor development and growth in nude mice that have been injected with suspended cells of the human colon cancer line, HCT-116.

Anti-human G₁₇ serum was obtained from rats immunized against hG₁₇(6)-DT. Normal rat serum was obtained from nonimmunized rats. Immunoglobulin fractions of the anti-G₁₇ serum and the normal rat serum were prepared by affinity chromatography, using Protein A Sepharose. Both immunoglobulin preparations were adjusted to a final protein concentration of 1 mg/ml in PBS. Measured by RIA, the G₁₇-direct antigen binding capacity (ABC) of the anti-human G₁₇ immunoglobulin preparation was 30 pg/ul. The normal rat immunoglobulin preparation had no anti-human G₁₇ activity.

HCT-116 cells were grown *in vitro* in the presence of pentagastrin as described in Example 6. Cells in the log phase of growth were collected from *in vitro* culture, washed by centrifugation in PBS, and resuspended to 10 cells/ml. Viability was assessed by trypan blue exclusion.

Each of 20 nude mice (NIH strain) were injected subcutaneously on their dorsal side near the right shoulder with a single bolus of 5×10^6 cells.

Two days after the injection of HCT-116 cells, the mice were randomly assigned to two groups of ten mice each. One group (Treated Group) was injected intraperitoneally with 0.5 mg per mouse of the anti-human G₁₇ immunoglobulin preparation. The other group (Control Group) was treated with 0.5 mg per mouse of the normal immunoglobulin fraction. Seven days later these treatments were repeated. These treatments resulted in sera G₁₇ binding capacities of 11.1 pg/ul in the anti-G₁₇ immunoglobulin treated mice and 0 in the normal immunoglobulin treated mice (See Table 5).

Two days after injections of the HCT-116 cells, all of the mice were started on daily injections of human G₁₇ for 16 consecutive days. A total daily dose of 51 ug of hormone per mouse was administered in 3 separate injections of 17 ug each, given at 4 hour intervals. The mice were bled on day 16 of the test to determine binding capacities by RIA.

Daily observations on the occurrence and growth of tumors were made visually and by palpation. On day 18 of the test, the tumors were measured by vernier caliper and the volume of each tumor estimated by the following formula: volume = (length X width²) ÷ 2 (Euhus et. al., 1986, J. Surg. Oncol. 31:229-234). On Day 18, the mice were euthanized and those mice without visually detectable tumors were dissected and further examined for tumors under a stereo microscope at 10X magnification.

As shown in Table 5, the anti-human G₁₇ immunoglobulin prevented tumors from developing in six out of ten mice in the Treated Group. Only one normal immunoglobulin-treated mouse failed to develop a tumor over the course of the test. In the four anti-G₁₇ treated animals that developed tumors, the mean tumor volume was reduced greater than four-fold compared to the tumors that developed in the mice treated with normal rat immunoglobulin. The Results demonstrate that treatment with anti-human G₁₇ immunoglobulin inhibits the development and growth of HCT-116 tumors in nude mice.

Table 5

	Treatment	
	Normal Rat Immunoglobulin	Anti-G ₁₇ Immunoglobulin
Mean Sera Anti-G ₁₇ ABC Titer (Range)	0.0 pg/ul	11.1 pg/ul (7.6-14.8)
Number of Mice Developing Tumors	9	4
Mean Tumor Volume ± S.E.	21.2 ± 11.2 mm	4.7 ± 2.8 mm

EXAMPLE 9

This test demonstrates that antibody - mediated immunity to G₁₇ can be selectively and safely reversed by the infection of peptide capable of binding to the antibody.

Six rats that were twice previously immunized with human gastrin G₁₇(6)-DT immunogen and which exhibited anti-human gastrin antigen binding capacity of 17-34 pg/ul of serum were prepared for the

standard stomach perfusion procedure. To demonstrate that each immunized rat was able to inhibit the acid secretion stimulatory activity of human G₁₇, a standard dose of approximately 120 ng of G₁₇ hormone was administered to each rat. After measuring the human gastrin G₁₇ stimulated acid output in response to the 120 ng of G₁₇, a second dose of human gastrin G₁₇, of 2.5 ug of hormone at twenty times the first dose was given to each rat and the gastrin stimulated acid output was measured again.

Two of the rats were then euthanized and their kidneys removed for sectioning and examination for deposition of complexes of gastrin and anti-gastrin antibody. Two of the remaining rats were then euthanized and their kidneys removed for sectioning. The remaining two rats were given 250 ug of human gastrin G₁₇, were followed for gastrin stimulated acid secretion, and then were euthanized and their kidneys were removed for sectioning.

Rapid reversal of human gastrin G₁₇ neutralizing activity of the anti-sera of human gastrin G₁₇ immunized rats is demonstrated in Figure 9. As expected from our previous perfusion assays, administration of approximately 120 ng of human gastrin G₁₇ resulted in an 87% mean inhibition of the expected gastrin stimulated acid secretion of all the rats. Challenge with 2.5 ug of human gastrin G₁₇ resulted in a gastrin stimulated acid response identical to that seen in non-immunized rats. Challenge with 25 ug or 250 ug of human gastrin G₁₇ resulted in an exaggerated acid secretion response. Kidney sections taken from these rats treated with large amounts of human gastrin G₁₇ were all negative for formation and deposition of immune complexes.

Treatment of rats immunized with human gastrin G₁₇ with only 2.5 ug of human gastrin G₁₇ immediately reversed the neutralization of gastrin stimulated acid secretion that was observed in these rats when they were first challenged with 120 ng of human gastrin G₁₇. Based on the antigen binding capacities of these rats, which exhibited a range of 17-34 pg/ul and a mean of 24 pg/ul, 2.5 ug of G₁₇ is at least a four-fold excess of hormone injected over the total antigen binding capacity of the rat's serum. Such a small amount of hormone, if given in the same proportion based on body weight to humans would amount to a range of only 500-700 ug of human G₁₇.

Preferably the antibody neutralizing by infusion would utilize a monovalent neutralizing molecule that bears the gastrin epitope but which does not itself induce acid secretion (e.g., by changing the C-terminal end of G₁₇).

The antibody neutralizer will not prevent a renewed production of anti-gastrin antibodies, the duration of which is determined by the conditions of immunization. However, it will neutralize the antibodies as they are produced. In practice, it may be necessary in most cases to provide for neutralization of antibodies synthesized after the initial dose of neutralizing compound is administered. This could be accomplished by means of additional infusions or, preferably, through the administration of the neutralizer in a sustained release compound or device. Although such administration would continue until the synthesis of anti-gastrin antibodies ceases, the quantity of antibody to be neutralized would be significantly less than that eliminated by the first administration of neutralizer. Consequently, the dose/frequency of neutralizer administrations would be diminished as antibody production subsides.

This invention and its preferred embodiments have been described in detail. It will be appreciated that those skilled in the art, upon consideration of this disclosure, may make modifications and improvements within the scope of this invention.

Claims

1. Immunogens comprising an immunogenic carrier conjugated to a peptide residue selected from the group of peptide residues which comprise fragments of the N-terminal amino acid sequence of heptadecagastrin ("G₁₇") up to amino acid residue number 12, and peptide residues, which comprise fragments of the N-terminal amino acid sequence of tetrateseracontagatrin ("G₃₄") up to amino acid residue number 22, for use in the preparation of pharmaceutical compositions useful in the treatment of gastrin-induced disorders.

2. Immunogens for use in the preparation of pharmaceutical compositions according to claim 1, wherein the amino acid sequence of the peptide residue differs by one or more amino acids from the natural fragments.

3. Immunogens for use in the preparation of pharmaceutical compositions according to claim 1, wherein the peptide residue comprises the fragment coupled to a spacer peptide chain.

4. Immunogens for use in the preparation of pharmaceutical compositions according to claim 3, wherein the spacer peptide chain is Arg-Pro-Pro-Pro-Cys-.

5. Immunogens for use in the preparation of pharmaceutical compositions according to claim 1, wherein

the peptide residue is selected from the group of peptide residues which comprise fragments of the N-terminal amino acid sequence of G₁₇, which are not contained in the N-terminal amino acid sequence of G₃₄.

6. Immunogens for use in the preparation of pharmaceutical compositions according claim 5, wherein
the peptide residue comprises one of the following amino acid sequences: pGlu-Gly-Pro-Trp-Leu-Glu-Glu-
Glu-Glu-Glu-Ala-Tyr, pGlu-Gly-Pro-Trp-Leu-Glu-, pGlu-Gly-Pro-Trp-Leu-Glu-Glu-, pGlu-Gly-Pro-Trp-Leu-Glu-
Glu-Glu-, pGlu-Gly-Pro-Trp-Leu-, pGlu-Gly-Pro-Trp-, pGlu-pGly-Pro-Trp-Leu-Glu-Glu-Glu-Glu-Glu-Ala, pGlu-
Gly-Pro-Trp-Leu-Glu-Glu-Glu-Glu-, pGlu-Gly-Pro-Trp-Leu-Glu-Glu-Glu-

7. Immunogens for use in the preparation of pharmaceutical compositions according to claim 5, wherein
10 the peptide residue comprises the amino acid sequence pGlu-Gly-Pro-Trp-Leu-Glu-.

8. Immunogens for use in the preparation of pharmaceutical compositions according to claim 6, wherein the peptide residue is selected from the group of the following peptide residues: pGlu-Gly-Pro-Trp-Leu-Glu-Arg-Pro-Pro-Pro-Pro-Cys-, pGlu-Gly-Pro-Trp-Leu-Arg-Pro-pro-Pro-Pro-Cys-, pGlu-Gly-Pro-Trp-Arg-Pro-Pro-Pro-Pro-Cys-, pGlu-Gly-Pro-Trp-Leu-Glu-Glu-Lys-Arg-Pro-Pro-Pro-Pro-Lys-.

15 9. Immunogens for use in the preparation of pharmaceutical compositions according to claim 1, wherein the peptide residue is selected from the group of peptide residues which comprise fragments of the N-terminal amino acid sequence G₃₄, which are not contained in the N-terminal amino acid sequence of G₁₇.

10. Immunogens for use in the preparation of pharmaceutical compositions according to claim 9, wherein the peptide residue comprises one of the following amino acid sequences: pGlu-Leu-Gly-Pro-Gln-Gly-Pro-Pro-His-Leu-Val-Ala-Asp-, pGlu-Leu-Gly-Pro-Gln-Gly-Pro-Pro-His-Leu-Val-Ala-Asp-Pro-Ser-Lys-Lys-Gln-Gly-Pro-Trp-Leu-, pGlu-Leu-Gly-Pro-Gln-Gly-, pGlu-Leu-Gly-Pro-Gln-Gly-Pro-Pro-His-, pGlu-Leu-Gly-Pro-Gln-Gly-Pro-Pro-His-Leu-Val-, pGlu-Leu-Gly-Pro-Gln-Gly-Pro-Pro-His-Leu-Val-Ala-Asp-Pro-Ser-Lys-Lys-Gln-Gly-Pro-Trp-, pGlu-Leu-Gly-Pro-Gln-Gly-Pro-Pro-His-Leu-Val-Ala-Asp-Pro-Ser-Lys-Lys-Gln-Gly-Pro-,

25 11. Immunogens for use in the preparation of pharmaceutical compositions according to claim 10, wherein the peptide residue comprises the amino acid sequence PGlu-Leu-Gly-Pro-Gln-Gly-.

12. Immunogens for use in the preparation of pharmaceutical compositions according to claim 10, wherein the peptide residue is: pGlu-Leu-Gly-Pro-Gln-Gly-Arg-Pro-Pro-Pro-Cys-.

13. Immunogens for use in the preparation of pharmaceutical compositions comprising an immunogenic carrier coupled to a peptide residue wherein the peptide residue comprises a fragment of the amino acid sequence of G₃₄ which is also contained in the amino acid sequence of G₁₇.

14. Immunogens for use in the preparation of pharmaceutical compositions according to claim 13, wherein the peptide residue has the amino acid sequence Asp-Pro-Ser-Lys-Lys-Gln-Gly-Pro-Trp-Leu-Pro-Pro-Pro-Cys, Asp-Pro-Ser-Lys-Lys-Gln-Gly-Pro-Trp-Leu, or Asp-Pro-Ser-Lys-Lys-Glu-Pro-Trp-Leu.

35 15. Immunogens for use in the preparation of pharmaceutical compositions according to claim 1, wherein the immunogenic carrier is selected from the group of diphtheria toxoid, tetanus toxoid, keyhole limpet hemocyanin, bovine serum albumin and fragments thereof.

17. Immunogens for use in the preparation of pharmaceutical compositions according to claim 1, wherein the peptide residues are bound on their C-terminal ends to a polymer consisting of crosslinked peptide units.

18. Immunogens for use in the preparation of pharmaceutical compositions according to claim 17, wherein the peptide residues are bound to an amino acid sequence comprising free amino groups which are crosslinked with glutaraldehyde.

19. A pharmaceutical composition comprising an immunogen comprising an immunogenic carrier conjugated to a peptide residue selected from the group of peptide residues which comprise fragments of the N-terminal amino acid sequence of heptadecagastrin ("G₁₇") up to amino acid residue number 12, and peptide residues, which comprise fragments of the N-terminal amino acid sequence of tetrateseracontagastrin ("G₃₄") up to amino acid residue number 22, and pharmaceutically acceptable carriers and/or adjuvants.

50 20. A pharmaceutical composition according to claim 19, wherein the amino acid sequence of the fragment portion of the peptide residue differs by one or more amino acids from the natural fragment.

21. A pharmaceutical composition according to claim 19, wherein the peptide residue comprises the fragment coupled to a spacer peptide chain.

55 22. A pharmaceutical composition according to claim 21, wherein the spacer peptide chain is Arg-Pro-Pro-Pro-Pro-Cys-.

23. A pharmaceutical composition according to claim 19, wherein the peptide residue is selected from the group of peptide residues which comprise fragments of the N-terminal amino sequence of G₁₇, which are not contained in the N-terminal amino acid sequence of G₃₄.

24. A pharmaceutical composition according to claim 22, wherein the peptide residue comprises one of the following amino acid sequences: pGlu-Gly-Pro-Trp-Leu-Glu-Glu-Glu-Glu-Ala-Tyr-, pGlu-Gly-Pro-Trp-Leu-Glu-, pGlu-Gly-Pro-Trp-Leu-Glu-Glu-, pGlu-Gly-Pro-Trp-Leu-Glu-Glu-Glu-, pGlu-Gly-Pro-Trp-Leu-, pGlu-Gly-Pro-Trp-

25. A pharmaceutical composition according to claim 21, wherein the peptide residue is selected from the group of the following peptide residues: pGlu-Gly-Pro-Trp-Leu-Glu-Arg-Pro-Pro-Pro-Pro-Cys-, pGlu-Gly-Pro-Trp-Leu-Arg-Pro-Pro-Pro-Pro-Cys-, pGlu-Gly-Pro-Trp-Arg-Pro-Pro-Pro-Pro-Cys-, pGlu-Arg-Pro-Pro-Leu-Glu-Arg-Pro-Pro-Pro-Pro-Cys-.

26. A pharmaceutical composition according to claim 19, wherein the peptide residue is selected from the group of peptide residues which comprise fragments of the N-terminal amino acid sequence of G₃₄, which are not contained in the N-terminal amino acid sequence of G₁₇.

27. A pharmaceutical composition according to claim 26, wherein the peptide residue comprises one of the following amino acid sequences: pGlu-Leu-Gly-Pro-Gln-Gly-Pro-Pro-His-Leu-Val-Ala-Asp-Pro-Ser-Lys-Lys-Gln-Gly-Pro-Trp-Leu-, pGlu-Leu-Gly-Pro-Gln-Gly-, pGlu-Leu-Gly-Pro-Gln-Gly-Pro-Pro-His-, pGlu-Leu-Gly-Pro-Gln-Gly-Arg-Pro-Pro-Pro-Pro-Cys-.

28. A pharmaceutical composition comprising an immunogen comprising an immunogenic carrier conjugated to a peptide residue, wherein the peptide residue comprises a fragment of the amino acid sequence of G₃₄ which is also contained in the amino acid sequence of G₁₇.

29. A pharmaceutical compositions according to claim 28, wherein the peptide residue has the amino acid sequence Asp-Pro-Ser-Lys-Lys-Gln-Gly-Pro-Trp-Leu-Pro-Pro-Pro-Pro-Cys-, Asp-Pro-Ser-Lys-Lys-Gln-Gly-Pro-Trp-Leu-, or Asp-Pro-Ser-Lys-Lys-Glu-Pro-Trp-Leu-.

30. A pharmaceutical composition according to claim 19, wherein the immunogenic carrier is selected from the group of diphtheria toxoid, tetanus toxoid, keyhole limpet hemocyanin, bovine serum albumin and fragments thereof.

31. A pharmaceutical composition according to claim 19, wherein the peptide residues are bound on their C-terminal ends to a polymer consisting of crosslinked peptide units.

32. A pharmaceutical composition according to claim 31, wherein the peptide residues are bound to an amino acid sequence comprising free amino groups which are crosslinked with glutaraldehyde.

33. Antibodies against gastrin hormone which selectively bind with one or both of G₁₇ and G₃₄ for use in the preparation of pharmaceutical compositions useful in the treatment of gastrin induced disorders.

34. Antibodies for use in the preparation of pharmaceutical compositions according to claim 33 which are generated from immunogens as defined in claim 1.

35. Pharmaceutical compositions comprising antibodies against gastrin hormone which selectively bind with one or both of G₁₇ and G₃₄ and pharmaceutically acceptable carriers and/or adjuvants.

36. Peptides which are void of gastrin activity, which comprise fragments of the N-terminal amino acid sequence of heptadecagastrin ("G₁₇") up to amino acid residue number 12 or which comprise fragments of the N-terminal amino acid sequence of tetrateseracontagastrin ("G₃₄") up to amino acid residue number 22, and which are capable of binding to gastrin antibodies as defined in claim 33 for use in the preparation of pharmaceutical compositions useful for reversing immunity to one or both of G₁₇ or G₃₄ mediated by said antibodies.

37. An immunogen comprising of a peptide selected from the group consisting of pGlu-Gly-Pro-Trp-Leu-Glu-Glu-Glu-, pGlu-Gly-Pro-Trp-Leu-Glu-Glu-, pGlu-Gly-Pro-Trp-Leu-Glu-, pGlu-Gly-Pro-Trp-Leu-, or pGlu-Gly-Pro-Trp coupled to an immunogenic carrier.

38. An immunogen comprising a peptide selected from the group consisting of a peptide pGlu-Leu-Gly-Pro-Gln-Gly-, Glu-Leu-Gly-Pro-Gln-Gly-Pro-Pro-His or pGlu-Leu-Gly-Pro-Gln-Gly-Arg-Pro-Pro-Pro-Cys coupled to an immunogenic carrier.

39. An immunogen comprising the peptide Asp-Pro-Ser-Lys-Lys-Gln-Pro-Trp-Leu coupled to an immunogenic carrier.

40. The immunogen of claim 37, 38, and 39 wherein the immunogenic carrier is coupled to diphtheria toxoid, tetanus toxoid, keyhole limpet hemocyanin or bovine serum albumin.

41. A peptide fragment comprising pGlu-Gly-Pro-Trp-Leu-Glu-Glu-Glu-, pGlu-Gly-Pro-Trp-Leu-Glu-Glu-, pGlu-Gly-Pro-Trp-Leu-Glu-, pGlu-Gly-Pro-Trp-Leu-, or pGlu-Gly-Pro-Trp.

42. A peptide fragment comprising pGlu-Leu-Gly-Pro-Gln-Gly-, pGlu-Leu-Gly-Pro-Gln-Gly-Pro-Pro-His or pGlu-Leu-Gly-Pro-Gln-Gly-Arg-Pro-Pro-Pro-Cys.

43. The peptide fragment Asp-Pro-Ser-Lys-Lys-Gln-Pro-Trp-Leu.

44. The use of a peptide of claim 41, 42 or 43 for the preparation of a pharmaceutically acceptable composition for the treatment of diseases caused by the activity of the hormone gastrin.

45. A pharmaceutical composition comprising a peptide fragment of claim 41, 42 or 43.

46. The use of a peptide of claim 41, 42 or 43 for the preparation of a pharmaceutically acceptable composition to the neutralizing anti-gastrin antibodies in vivo.

5

10

15

20

25

30

35

40

45

50

55




FIGURE 1**Figure 1. G17 with Control Rat Serum****Figure 2. G17 with Rat anti-G17 Serum**

Figure 3


Stomach Acid Output in Response to
Injection of G17 and Pentagastrin (pG)
in Rat Actively Immunized Against G17



Arrows: Time of G17 or pG injection.
Acid Secretion peaks are numbered 1,2,3.

Figure 4


Stomach Acid Output in Response to
Injection of G17 and Pentagastrin (pG)
in Rat Actively Immunized Against G17



Arrows: Time of G17 or pG Injection.
Acid secretion peaks are numbered 1,2,3.

Figure 5


**Stomach Acid Output in Response to
Injection of G17 and Pentagastrin (pG)
in Control Rat**



**Arrows: Time of G17 or pG Injection.
Acid secretion peaks are numbered 1,2,3.**

Figure 6


Stomach Acid Output in Response to
Injection of G17 and G34
in Rat Actively Immunized Against G17



Arrows: Time of G17 or G34 injection.
Acid Secretion peaks are numbered 1,2,3.

Figure 7


Stomach Acid Output in Response to
Injection of G17 and G34
in Control Rat



Arrows: Time of G17 or G34 injection.
Acid Secretion peaks are numbered 1,2,3.


Figure 8

Mouse Antibody Responses to G-17 After Immunization with Polymerized Synthetic Peptides



Binding capacities determined by RIA,
using $(^{125}\text{I})\text{G-17}$. Values Plotted =
Mean (- Bknd) and s.e; n = 5.


**Figure 9 Total Quantity of Acid
Secreted by hG17-Immune and Non-Immune
Rats in Response to Graded Doses of hG17**




n=6 for rats injected with .12 & 2.5ug.
n=2 for rats injected with 25 and 250ug.

FIGURE 10

Stimulation of the Growth of Human Colon Cancer Cell Line HCT 116 By Pentagastrin




**Figure 11 : Effect of Anti-G17 Ab on
the Growth of Colon Cancer Implants in
Nude Mice Infused with G17**



Gp I: Injected with Rat Anti-G17 Ab.
Gp II: Injected with Normal Rat Ab.

**Figure 12 : Effect of Anti-G17 Ab on
the Growth of Colon Cancer Implants in
Nude Mice**



Group I: Injected with G17 and Anti-G17.
Group III: Injected with saline (no G17)
and not injected with Ab.



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number:

0 380 230 A3

12

EUROPEAN PATENT APPLICATION

21 Application number: 90300456.2

51 Int. Cl.⁵: **C07K 15/00, A61K 39/385,
A61K 37/24, C07K 99/30**

22 Date of filing: 17.01.90

30 Priority: 24.01.89 US 301353
12.05.89 US 351193

43 Date of publication of application:
01.08.90 Bulletin 90/31

84 Designated Contracting States:
AT BE CH DE DK ES FR GB GR IT LI LU NL SE

88 Date of deferred publication of the search report:
26.06.91 Bulletin 91/26

71 Applicant: **APHTON CORPORATION**
P.O. Box 1049
Woodland, CA 95695(US)

72 Inventor: **Gevas, Philip C.**
487-A Portlock Road
Honolulu, Hawaii 96825(US)
Inventor: **Grimes, Stephen**
1400 Cypress Lane
Davis, California 95616(US)
Inventor: **Karr, Stephen L., Jr.**
624 Villanova Drive
Davis, California 95616(US)
Inventor: **Littenberg, Richard L.**
70 Kai One Place
Kai Lua, Hawaii 96734(US)


74 Representative: **Lucas, Brian Ronald**
Lucas, George & Co. 135 Westhall Road
Warlingham Surrey CR3 9HJ(GB)

54 Immunogenic compositions against gastrin peptides.

57 Immunogenic compositions useful for the treatment of ulcers or tumors whose growth is dependent on or stimulated by gastrin hormones are disclosed. The immunogenic compositions induce antibodies in a subject which selectively neutralize the specific hormones. Pharmaceutical compositions comprising effective amounts of the immunogenic compositions and methods of treatment using the compositions are disclosed. A method of reversing the inventive treatments by neutralizing the antibodies induced in vivo is also disclosed.

Figure 3

Stomach Acid Output in Response to
Injection of G17 and Pentagastrin (pG)
in Rat Actively Immunized Against G17



Arrows: Time of G17 or pG injection.
Acid Secretion peaks are numbered 1,2,3.

EP 0 380 230 A3



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number

EP 90 30 0456

Page 1

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
Y	BIOLOGICAL ABSTRACT Nb 71058738 CHOUDHURY A.M. et al : " Amino terminal sequence of human big gastrin sequence synthetic and immunochemical studies." & HOPPE-SEYLER'S Z. PHYSIOL. CHEM. 361, (11), 1980, 1719-1734.	1, 9, 13, 19, 26, 28, 36	C07K15/00 A61K39/385 A61K37/24 C07K99/30
X	---	33-35	
Y	MEDLINE ABSTRACT Nb 81200269 DOCKRAY G.J. : " Immunochemical studies on big gastrin using NH-2 terminal specific antisera." & REGUL. PEP., december 1980, 1 (3), 169-186.	1, 9, 13, 19, 26, 28, 36	
X	---	33-35	
X	BIOLOGICAL ABSTRACT Nb 63037293 DOCKRAY G.J. et al: " Hepta deca peptide gastrin measurement in blood by specific radio immunoassay. & GASTROENTEROLOGY, 1976, 71 (6), 971-77.	33-35	
A	CHEMICAL ABSTRACTS, vol. 109, no. 9, 29 August 1988 Columbus, Ohio, USA POWER D. M. et al: "A novel gastrin-processing pathway in mammalian antrum." page 113; column 1; ref. no. 67341 Z & BIOCHIM. BIOPHYS. ACTA 1988, 954(2), 141-7 * abstract *	1-46	TECHNICAL FIELDS SEARCHED (Int. Cl.5)
A	CHEMICAL ABSTRACTS, vol. 92, no. 21, 26 May 1980 Columbus, Ohio, USA WENDLBERGER G. et al: "The syntheses of human big gastrin I and its 32-lucifine analog. 2. Preparation of fragments 9-14 and 1-8." page 722; column 2; ref. no. 198749 S & MONATSH. CHEM. 1979, 110(6), 1317-30. * abstract *	1-46	C07K A61K
A	US-A-4384995 (STEVENS V.C.) * column 3 *	1-46	
D	& US-A-4302386		
The present search report has been drawn up for all claims			
Place of search BERLIN		Date of completion of the search 18 APRIL 1991	Examiner AVEDIKIAN P. F.
CATEGORY OF CITED DOCUMENTS			
X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document I : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503 03.82 (P0401)